

# Assignment 0

- State TRUE or FALSE giving proper justification for each of the following statements.
  - There exists an unbounded subset  $A$  of  $\mathbb{R}$  such that  $m^*(A) = 5$ .
  - There exists an open subset  $A$  of  $\mathbb{R}$  such that  $[\frac{1}{2}, \frac{3}{4}] \subset A$  and  $m(A) = \frac{1}{4}$ .
  - There exists an open subset  $A$  of  $\mathbb{R}$  such that  $m(A) < \frac{1}{5}$  but  $A \cap (a, b) \neq \emptyset$  for all  $a, b \in \mathbb{R}$  with  $a < b$ .
  - If  $A$  and  $B$  are open subsets of  $\mathbb{R}$  such that  $A \subsetneq B$ , then it is necessary that  $m(A) < m(B)$ .
  - If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous a.e. on  $\mathbb{R}$ , then there must exist a continuous function  $g : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f = g$  a.e. on  $\mathbb{R}$ .
  - If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and if  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that  $f = g$  a.e. on  $\mathbb{R}$ , then  $f$  must be continuous a.e. on  $\mathbb{R}$ .
- Let  $f : [0, 2) \rightarrow \mathbb{R}$  be defined by  $f(x) = \begin{cases} x^2 & \text{if } 0 \leq x \leq 1, \\ 3 - x & \text{if } 1 < x < 2. \end{cases}$   
Find  $m^*(A)$ , where  $A = f^{-1}((\frac{9}{16}, \frac{5}{4})) = \{x \in [0, 2) : f(x) \in (\frac{9}{16}, \frac{5}{4})\}$ .
- Let  $B \subset A \subset \mathbb{R}$  such that  $m^*(B) = 0$ . Show that  $m^*(A \setminus B) = m^*(A)$ .
- Let  $A \subset \mathbb{R}$  such that  $m^*(A) > 0$ . Show that there exists  $B \subset A$  such that  $B$  is bounded and  $m^*(B) > 0$ .
- If  $A \subset \mathbb{R}$ , then show that  $m^*(A) = \inf\{m(G) : A \subset G, G \text{ is an open set in } \mathbb{R}\}$ .
- Let  $E = \{x \in [0, 1] : \text{The decimal representation of } x \text{ does not contain the digit } 5\}$ . Show that  $m(E) = 0$ .
- If  $G$  is a nonempty open subset of  $\mathbb{R}$ , then show that  $m(G) > 0$ .
- Show that a subset  $E$  of  $\mathbb{R}$  is Lebesgue measurable iff  $m^*(I) = m^*(I \cap E) + m^*(I \setminus E)$  for every bounded open interval  $I$  of  $\mathbb{R}$ .
- Let  $A, B \subset \mathbb{R}$  such that  $m^*(A) = 0$  and  $A \cup B$  is Lebesgue measurable. Show that  $B$  is Lebesgue measurable.
- Let  $A, B \subset \mathbb{R}$  such that  $A$  is Lebesgue measurable and  $m^*(A \Delta B) = 0$ . Show that  $B$  is Lebesgue measurable.
- Let  $A \subset \mathbb{R}$  such that  $A \cap B$  is Lebesgue measurable for every bounded subset  $B$  of  $\mathbb{R}$ . Show that  $A$  is Lebesgue measurable.
- If  $E$  is a Lebesgue measurable subset of  $\mathbb{R}$  and if  $x \in \mathbb{R}$ , then show that  $E + x$  is Lebesgue measurable.
- Let  $A$  be a countable subset of  $\mathbb{R}$  and let  $B \subset \mathbb{R}$  such that  $m^*(B) = 0$ . Show that  $m^*(A+B) = 0$ .
- Let  $I$  and  $J$  be disjoint open intervals in  $\mathbb{R}$  and let  $A \subset I, B \subset J$ . Show that  $m^*(A \cup B) = m^*(A) + m^*(B)$ .

15. Let  $A \subset [0, 1]$  be Lebesgue measurable with  $m(A) = 1$ . If  $B \subset [0, 1]$ , then show that  $m^*(A \cap B) = m^*(B)$ .
16. Let  $E_i \subset (0, 1)$  ( $i = 1, \dots, n$ ) be Lebesgue measurable sets such that  $\sum_{i=1}^n m(E_i) > n - 1$ . Show that  $m(\cap_{i=1}^n E_i) > 0$ .
17. If  $A \subset \mathbb{R}$ , then show that there exists a Lebesgue measurable subset  $E$  of  $\mathbb{R}$  such that  $m^*(A) = m(E)$ .
18. Let  $A \subset \mathbb{R}$  such that  $m^*(A) > 0$ . Show that there exist  $x, y \in A$  such that  $x - y \in \mathbb{R} \setminus \mathbb{Q}$ .
19. Show that the Borel  $\sigma$ -algebra on  $\mathbb{R}$  is generated by the class  $\{(-\infty, x] : x \in \mathbb{Q}\}$ .
20. Let  $E \subset \mathbb{R}$  and let  $\alpha \in \mathbb{R}$ . If  $\alpha E = \{\alpha x : x \in E\}$ , then show that  $m^*(\alpha E) = |\alpha|m^*(E)$ . Also, show that if  $E$  is Lebesgue measurable, then  $\alpha E$  is Lebesgue measurable.
21. Let  $A \subset \mathbb{R}$  such that  $m^*(A) = 0$ . Show that  $m^*(\{x^2 : x \in A\}) = 0$ .
22. Let  $A, B \subset \mathbb{R}$  such that  $A \cup B$  is Lebesgue measurable and  $m(A \cup B) = m^*(A) + m^*(B) < \infty$ . Show that both  $A$  and  $B$  are Lebesgue measurable.
23. If  $A \subset \mathbb{R}$ , then show that  $\chi_A$  is a Lebesgue measurable function iff  $A$  is a Lebesgue measurable set.
24. Let  $E$  be a Lebesgue measurable subset of  $\mathbb{R}$ . Show that  $f : E \rightarrow \mathbb{R}$  is Lebesgue measurable iff  $\{x \in E : f(x) > r\}$  is Lebesgue measurable for each  $r \in \mathbb{Q}$ .
25. Let  $E$  be a Lebesgue measurable subset of  $\mathbb{R}$  and let  $f : E \rightarrow \mathbb{R}$  be a Lebesgue measurable function. For each  $x \in E$ , let  $g(x) = \begin{cases} f(x) & \text{if } |f(x)| \leq 5, \\ 0 & \text{if } |f(x)| > 5. \end{cases}$  Show that  $g : E \rightarrow \mathbb{R}$  is Lebesgue measurable.
26. Let  $E$  be a Lebesgue measurable subset of  $\mathbb{R}$  and let  $f : E \rightarrow \mathbb{R}$  be a Lebesgue measurable function. For each  $x \in E$ , let  $g(x) = \begin{cases} 0 & \text{if } f(x) \in \mathbb{Q}, \\ 1 & \text{if } f(x) \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$  Show that  $g : E \rightarrow \mathbb{R}$  is Lebesgue measurable.
27. Let  $E$  be a Lebesgue measurable subset of  $\mathbb{R}$  and let  $f : E \rightarrow \mathbb{R}$  be a Lebesgue measurable function. If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is continuous, then show that  $g \circ f$  is Lebesgue measurable.
28. Does there exist a continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f = \chi_{[0,1]}$  a.e. on  $\mathbb{R}$ ? Justify.
29. Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function. Show that  $f' : [a, b] \rightarrow \mathbb{R}$  is Lebesgue measurable.
30. If  $E$  is a Lebesgue measurable subset of  $\mathbb{R}$  with  $m(E) < +\infty$  and if  $f(x) = m(E \cap (-\infty, x])$  for all  $x \in \mathbb{R}$ , then show that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous.

31. Let  $E$  be a Lebesgue measurable subset of  $\mathbb{R}$  with  $m(E) < \infty$ . Define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) = m\{E \cap (-\infty, x^2)\}$ . Show that  $f$  is differentiable at 0 and  $f'(0) = 0$ .

32. For each  $x \in [0, 1]$ , let  $f(x) = \begin{cases} \frac{1}{n} & \text{if } x = \frac{m}{n} \text{ for some } m, n \in \mathbb{N} \text{ with g.c.d.}(m, n) = 1, \\ 0 & \text{otherwise.} \end{cases}$

Evaluate the Lebesgue integral  $\int_{[0,1]} f$ .

33. For each  $x \in [0, 1]$ , let  $f(x) = \begin{cases} x^2 & \text{if } x = \frac{1}{2^n} \text{ for some } n \in \mathbb{N}, \\ x^3 & \text{if } x = \frac{1}{3^n} \text{ for some } n \in \mathbb{N}, \\ x^4 & \text{otherwise.} \end{cases}$

Evaluate the Lebesgue integral  $\int_{[0,1]} f$ .

34. Let  $f(x) = \begin{cases} \sin(\pi x) & \text{if } x \in [0, \frac{1}{2}] \setminus C, \\ \cos(\pi x) & \text{if } x \in (\frac{1}{2}, 1] \setminus C, \\ x^2 & \text{if } x \in C. \end{cases}$

( $C$  denotes the Cantor set.) Evaluate the Lebesgue integral  $\int_{[0,1]} f$ .

35. Evaluate the Lebesgue integral  $\int_{[0,\infty)} e^{-[x]} dx$ .

36. Let  $f(x) = \begin{cases} e^{[x]} & \text{if } x \in \mathbb{Q}, \\ e^{-[x]} & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$

Evaluate the Lebesgue integral  $\int_{(0,\infty)} f$ .

37. Let  $f(x) = \begin{cases} e^{|x|} & \text{if } x \in \mathbb{Q}, \\ e^{-|x|} & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$

Evaluate the Lebesgue integral  $\int_{\mathbb{R}} f$ .

38. Evaluate the Lebesgue integral  $\int_{(0,1]} \frac{1}{\sqrt[3]{x}} dx$ .

39. Let  $f(x) = \begin{cases} \frac{1}{\sqrt{x}} & \text{if } 0 < x \leq 1, \\ \frac{1}{x} & \text{if } x > 1. \end{cases}$

Evaluate the Lebesgue integral  $\int_{(0,\infty)} f$ .

40. Evaluate the following:

(a)  $\lim_{n \rightarrow \infty} \int_{-2}^2 \frac{x^{2n}}{1+x^{2n}} dx$

(b)  $\lim_{n \rightarrow \infty} \int_{[0,1]} \frac{1+nx}{(1+x)^n} dx$

(c)  $\int_0^1 \left( \sum_{n=1}^{\infty} \frac{x^n}{n} \right) dx$

(d)  $\lim_{n \rightarrow \infty} \int_1^{\infty} \frac{1}{1+x^{2n}} dx$

(e)  $\sum_{n=0}^{\infty} \int_0^1 \frac{x^2}{(1+x^2)^n} dx$